

EXTRACTION AND ANTIOXIDANT EVALUATION OF EXTRACT OF *Eysenhardtia polystachya* FOR COSMETIC APPLICATIONS

ID
EP_507

González-Reza, Ricardo Moisés¹, Zambrano Zaragoza, María de la Luz^{1*}, Quintanar Guerrero, David²

¹Laboratorio de Procesos de Transformación y Tecnologías Emergentes de Alimentos, Universidad Nacional Autónoma de México, FES-Cuautitlan, Cuautitlan Izcalli 54714, Estado de México, México

²Laboratorio de Investigación y Posgrado en Tecnología Farmacéutica, Universidad Nacional Autónoma de México, FES- Cuautitlan, Cuautitlan Izcalli 54745, Estado de México, México.

Introduction:

Currently, cosmetology is looking for alternatives and new trends for skincare and sun protection; therefore, finding sustainable and high-impact alternatives in various products' new formulations is necessary. For example, *Eysenhardtia polystachya* is a tree distributed in Mexico and Texas and locally known as "palo azul." The infusion of this tree's bark in water shows fluorescence, and its components have been traditionally used for different purposes such as antidiabetic, antioxidant, antisolar, and antimicrobial. However, extraction is often inefficient and influences the properties and functionality, particularly on antioxidant capacity. This tree gained attention due to its blue fluorescence because of its infusion in water and its biological properties, including antidiabetic, antibacterial, and antioxidant (Narvaez-Mastache et al., 2006; Gutiérrez & Báez, 2014). Among the components that have been found in *E. polystachya* extracts are aurons, flavonoids, isoflavonoids, and matalin. All flavonoids contain a 15-carbon skeleton; their central structure is a 2-phenylbenzopyranone, in which the three-carbon bridge between phenyl groups is commonly cyclized with oxygen. Other components of the plant extract are methylated flavonoids (Ferreira-García et al., 2017). The compounds responsible for the fluorescence of *E. Polystachya* are Coatline A and B, which represent a C- β -glucopyranosyl- α -hydroxy dihydrochalcone (Beltrami et al., 1980) and different isoflavonoids (Burns et al., 1984). These compounds are ionizable at basic pH values, and fluorescence is observed (Álvarez & Delgado, 1999; Valeur, 2001). The work aimed was to establish the effectiveness of ultrasound extraction of *E. polystachya* on the phenolic and antioxidant capacity of extract.

Results & Discussion:

The spectral scan (Figure 1) revealed two maximum absorbances at 284 and 325 nm corresponding to two superimposed chromophore groups characteristic of the extract at 240-280 nm for the band I and at 320-380 nm for band II, verifying the presence of isoflavones, flavones, and flavanols.




Figure 1. The spectral scan of *E. polystachya* extract.



The concentration of polyphenols obtained was 92.42 mg GAEq/mg of extract, and for flavonoids, 59.91 mg quercetin Eq/mg, the extract obtained is rich in polyphenols than 65% are flavonoids. These values are comparable to those reported by Gutiérrez & Báez (2014) for EP extracts. Phenolic compounds have an aromatic ring that carries one or more hydroxyl groups, and their structure can vary from a simple phenolic molecule to that of a complex high molecular weight polymer (Haminiuk et al., 2012)

For the antioxidant capacity, the following results were obtained: 3.29 ± 0.11 μmol EAA/mg (FRAP), 2.97 ± 0.27 μmol EAA/mg (DPPH), and 2.82 ± 0.31 μmol EAA/mg (ABTS), which confirms the high antioxidant capacity of the extract obtained, which is assigned directly to the flavonoid-rich fraction it contains. Gutiérrez & Báez (2014) report that the results of antioxidant capacity obtained by ABTS and DPPH for the EP extracts are comparable for inhibiting free radicals when BHT and ascorbic acid are used positive control, inhibiting almost 100% of the radical. At concentrations of 100 $\mu\text{g/mL}$ of EP extract.

Materials & Methods:

Obtaining the extract

E. Polystachya was acquired in the municipal market of Cuautitlán México. The fraction rich in flavonoids was extracted with an ethanol/water mixture in a 2:1 ratio assisted by ultrasound using an ultrasonic processor (Hielscher Ultrasonics gmbh, UP200Ht, Teltow, Germany) in 3 h doing the ultrasonication every 20 min with a duration of 5 min each at a temperature of 25 °C with 50 W of power. The mass/volume ratio of the product and extraction medium was 1:6. After extraction, the solution was evaporated at 6.7 kPa in a rotary evaporator (RV10, IKA® Wilmington, USA) at 40 °C. Finally, the remaining solution was dried in trays in a convective oven at 50 °C to obtain the solids extracted rich in flavonoids.



Conclusions:

It can be concluded that the extract rich in flavonoids obtained in the present experimental study presents a great alternative as an antioxidant photoprotective agent for cosmetic applications, in particular for antiaging and sun protection products.

Aknowledgments:

The authors also acknowledge the financial support for PAPIIT IN222420 projects IN222520 of DAGAPA-UNAM and Internal Project-FESC PIAPI 2060 and 2040.

References:

- Alvarez, L., Delgado, G., (1999). C- and O-glycosyl- α -hydroxydihydrochalcones from *Eysenhardtia polystachya*. *Phytochemistry*, 50: 681-687.
- Beltrami, E., De Bernardi, M., Fronza, G., Mellerio, G., Vidari, G., Vita-Finzi, P. (1980). Coatline A and B, two C-glycosyl- α -hydroxydihydrochalcones from *Eysenhardtia polystachya*. *Phytochemistry*, 21: 2931-2933.
- Benzie, I. F., & Strain, J. J. (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical biochemistry*, 239(1), 70-76.
- Brand-Williams, W., Cuvelier, M. E., Berset, C. (1995). Use of a free radical method to evaluate antioxidant activity. *LWT - Food Science and Technology*, 28(1), 25-30.
- Burns, D.T., Dalgarno, G., Gargan, E., Gargan, P.E., Grimshaw, J. (1984). An isoflavone and a coumestan from *Eysenhardtia polystachya*—Robert Boyle's fluorescent acid-base indicator. *Phytochemistry*, 23: 167-169.
- Chang, C. C., Yang, M. H., Wen, H. M., & Chern, J. C. (2002). Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of food and drug analysis*, 10(3).
- Ferreira-García, M.G., Hernandez-Martinez, A.R., Esparza, R., Molina, G.A., Rodriguez-Melgarejo, F., Jiménez, S., Estevez, M. (2017). Effects of extraction solvents on photoluminescent properties of *Eysenhardtia polystachya* and their potential usage as biomarker. *Materials Science and Engineering C*, 72, 42-52.
- Gutiérrez R.M.P., Baez E.G. (2014). Evaluation of antidiabetic, antioxidant and antiglycating activities of the *Eysenhardtia polystachya*. *Pharmacognosy Magazine*, 10: S404-S418.
- Haminiuk, C.W., Maciel, G.M., Plata-Oviedo, M.S., Peralta, R.M. (2012). Phenolic compounds in fruits - an overview. *International Journal of Food Science and Technology*, (47), 2023-2044.
- Narvaez-Mastache J.M., Garduño-Ramírez M.L., Alvarez L., Delgado G. (2006). Antihyperglycemic activity and chemical constituents of *Eysenhardtia platycarpa*. *Journal of Natural Products*, 69: 1687-1691.
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans, C. (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*, 26(9): 1231-1237.
- Singleton, V.L., Rossi, J.A.Jr., (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, 16, 144-158.
- Valeur, B. (2001). Fluorescent molecular sensors of ions and molecules. *Molecular Fluorescence: Principles and Applications*, 273-350.