

# Nanocomposite of layered double hydroxide and chitosan polymer for delivery of retinol and Pal-RGD

GS\_328

Kim, Tae Ho<sup>1</sup>; Kim, Soo Yeon<sup>1</sup>; Lee, Jong Hyeon<sup>1\*</sup>  
Park, Young Ah<sup>2,3</sup>; Kang, Yu Jin<sup>2</sup>; Ji, Hong Geun<sup>2</sup>

<sup>1</sup> Department of Chemistry, The Catholic University of Korea, Bucheon, 14662, Korea  
<sup>2</sup> H&A PharmaChem, R&D center, Bucheon, 14558, Korea  
<sup>3</sup> Interdisciplinary Program in BioCosmetics, Sungkyunkwan University, Suwon, 16419, Korea  
\* Corresponding author email: [jhlee305@catholic.ac.kr](mailto:jhlee305@catholic.ac.kr)

## Introduction:

Layered double hydroxide (LDH) has been considered a potential host material in the functional delivery system in many academic and industrial fields. However, natural LDH contains harmful heavy metals, and new LDH synthesis for LDH-polymer composites has low productivity. More importantly, the one-pot synthesis of LDH-polymer composite does not guarantee the uniform LDH and its well-dispersion in a polymer matrix. Additionally, it requires a mass washing step to remove reaction residues. In this study, we investigated the porous nanocomposite of eco-friendly LDH nanosheets and naturally abundant chitosan biopolymer for uptake and selective release of cosmetic niosome vesicles ranging from 50 to 200 nm. Niosome plays an essential role in delivering active ingredients to the skin, leading to improved effectiveness of cosmetics. Although niosome nanoparticles are nontoxic and biodegradable, they have significant disadvantages, including low stability, the possibility of breaking down when in contact with skin, and short lifetimes. Also, they have a limited ability to prevent degradation of active ingredients caused by the penetration of oxygen gas and ultraviolet light. We suggest the use biocompatible LDH and chitosan biopolymer nanocomposites as a stabilizing carrier for colloidal niosome nanoparticles.

## Results & Discussion:

## Materials & Methods:




Figure 2. Schematic representation for calcined LDH, chitosan solution, Niosome carrying retinol and pal-RGD. Preparing step and simple image of calcined LDH-chitosan polymer (RLDH-CS) Niosome composite.




Figure 3. (A) Characteristics of Retinol and pal-RGD (B) Nanoparticle Tracking Analysis image and graph of Niosome, (a) Retinol (b) pal-RGD  
Table 1. Peak analysis according to NTA result of Niosome

### I. Retinol




Figure 4. (A) Schematic representation for calcined LDH-chitosan polymer (RLDH-CS) Niosome composite manufacturing process. Niosome is carrying retinol. (B) XRD pattern of LDH, RLDH-W and RLDH-CS Niosome nanocomposite and broadened as compared with LDH and RLDH-W. (C) UV/vis absorption spectroscopy of LDH, chitosan polymer, niosome, RLDH-CS nanocomposite and RLDH-CS Niosome nanocomposite. Absorbance was measured using an integrating sphere.

### II. pal-RGD




Figure 5. (A) Schematic representation for calcined LDH-chitosan polymer (RLDH-CS) Niosome composite manufacturing process. Niosome is carrying pal-RGD. (B) XRD pattern of LDH, RLDH-W, pal-RGD and RLDH-CS Niosome nanocomposite and broadened as compared with LDH and RLDH-W. (C) FT-IR spectrum clearly showed the characteristic peaks for pal-RGD at 1365, 1566, 1634, 1757, 2851, 2918 cm<sup>-1</sup>

### III. Uptake & Release




Figure 6. (A) Niosome uptake by RLDH-CS (B) Niosome release from the RLDH-CS, at two solution pH value: 6 and 7

## Conclusions:

We have used a commercial LDH to obtain the LDH nanosheet by the calcination-reconstruction method in the 3% acetic acid solution of chitosan polymer. In FT-IR spectra, a broad band but dominant band for OH vibration around 3400 cm<sup>-1</sup> compared to pure chitosan and the partial shift of Mg(Al)-O band toward lower frequencies would support the chemical interaction between the LDH and chitosan. Considering the hydrolysis and deprotonation of calcined LDH, it would expect a grafting of chitosan polymer on the active LDH surface during the reconstruction. Additionally, the nanocomposite is very stable both in low pH solutions and in the recycling conditions, mainly due to the chemical bonding of LDH surface and chitosan polymer. The nanocomposite exhibited the excellent uptake and release behavior for retinol and pal-RGD incorporated niosome. Therefore, the result explains that the nanocomposite's capacity potential is mainly ascribed to the positive surface charge of LDH nanosheets and polymer network in the nanocomposite.

## Acknowledgments:

This research was supported by the Technology Innovation Program (or Industrial Strategic Technology Development Program (10077704, Hybrid composite based on LDH and MOF for improved skin permeability and On/Off controllable function of physiologically active substance) funded by the Ministry of Trade, Industry & Energy (MOTIE, Korea).

## References:

1. Yu, J., et al. (2017). "Preparation of two dimensional layered double hydroxide nanosheets and their applications." *Chemical Society Reviews* 46(19): 5950-5974.
2. Khan, A. I. and D. O'Hare (2002). "Intercalation chemistry of layered double hydroxides: recent developments and applications." *Journal of Materials Chemistry* 12(11): 3191-3198.
3. Newman, S. P. and W. Jones (1998). "Synthesis, characterization and applications of layered double hydroxides containing organic guests." *New Journal of Chemistry* 22(2): 105-115.
4. Al-Jamal, W. T. and K. Kostarelos (2011). "Liposomes: from a clinically established drug delivery system to a nanoparticle platform for theranostic nanomedicine." *Accounts of chemical research* 44(10): 1094-1104.
5. Khezri, K., et al. (2018). "Application of nanoparticles in percutaneous delivery of active ingredients in cosmetic preparations." *Biomedicine & Pharmacotherapy* 106: 1499-1505.
6. Akbarzadeh, A., et al. (2013). "Liposome: classification, preparation, and applications." *Nanoscale research letters* 8(1): 1-9.
7. Kim, J., et al. (2021). "Porous nanocomposite of layered double hydroxide nanosheet and chitosan biopolymer for cosmetic carrier application." *Applied Clay Science* 205: 106067.